

*Hannu Vartiainen — Klaus Kultti*  
**Multilateral Non-Cooperative  
Bargaining  
in a General Utility Space**

**Aboa Centre for Economics**

Discussion Paper No. 19  
Turku 2007



Copyright © Author(s)

ISSN 1796-3133

Turun kauppakorkeakoulun monistamo  
Turku 2007

*Hannu Vartiainen – Klaus Kultti*  
**Multilateral Non-Cooperative Bargaining  
in a General Utility Space**

**Aboa Centre for Economics**  
Discussion Paper No. 19  
September 2007

**ABSTRACT**

We consider an  $n$ -player bargaining problem where the utility possibility set is compact, convex, and strictly comprehensive. We show that a stationary subgame perfect Nash equilibrium exists, and that, if the Pareto surface is differentiable, all such equilibria converge to the Nash bargaining solution as the length of a time period between offers goes to zero. Without the differentiability assumption, convergence need not hold.

JEL Classification: C7, D7

Keywords: multilateral, bargaining, general utility set

## **Contact information**

Klaus Kultti, University of Helsinki and Hannu Vartiainen, Turku School of Economics.

## **Acknowledgements**

We thank Geir Asheim and Hannu Salonen for comments and discussions.

# 1 Introduction

One of the most celebrated observations in the bargaining literature is by Binmore, Rubinstein and Wolinsky (1985), who show that the unique subgame perfect equilibrium outcome of the two-player alternating offers bargaining game à la Rubinstein (1982) converges to the Nash bargaining solution when frictions vanish, i.e., when the length of time period that it takes to make offers and counter offers goes to zero.

We study how far these results extend to a multiplayer setting. To do this, we employ the natural multiplayer version of the alternating offers bargaining game by Binmore (1985), Herreró (1985), and Shaked (as reported by Osborne and Rubinstein, 1990).<sup>1</sup>

Vastness of equilibria is a well known problem of multiplayer bargaining games. To circumvent the problem, we shall focus on equilibria in *stationary* strategies. It is known that under stationarity restriction convergence does hold (see Sutton, 1986) when bargaining concerns linear, one dimensional cake.<sup>2</sup> Our question is whether convergence holds if we relax the restrictions on the underlying physical structure.

While in the two player case the underlying physical structure is immaterial, in the multiplayer case it is not. The one-cake restriction removes a large portion of interesting trade-offs between players over which bargaining could take place. In fact, we prove via an example that convergence of stationary equilibria to the Nash solution need not hold when the utility domain is only asked to meet the more liberal assumptions made in the cooperative bargaining literature, i.e. that the payoffs are drawn from a compact, convex and comprehensive utility possibility set.<sup>3</sup> Such more general utility set can be generated e.g. by multiplicity of goods or consumption externalities. Our aim is to delineate conditions under which convergence does take place in such a utility set.

We start by showing that a stationary subgame perfect equilibrium exists in any compact, convex, and comprehensive utility possibility set. To our knowledge, this is the most general existence result concerning stationary bargaining equilibria. Our main result is that by slightly restricting the utility space the convergence result can be saved: *all* stationary equilibria converge to the Nash bargaining solution if the Pareto surface is *differentiable*.<sup>4</sup>

A distinctive feature of our analysis is - since no specific correspondence

---

<sup>1</sup>Players are ordered into a circle. A player (say 1) proposes an outcome. If all other players accepts the proposal, then it is implemented. Otherwise, the next player (say 2) in the order makes an offer. The play continues this way until an offer is accepted.

<sup>2</sup>Chatterjee and Sabourian (2000) show that the stationarity restriction is without loss of generality if players are complexity averse.

<sup>3</sup>Thomson and Lensberg (1989) is an authoritative reference.

<sup>4</sup>We thank an associate editor for suggesting this weakening of our original restriction.

need be determined between payoffs and the physical environment - that a closed form representations of strategies and payoffs cannot be used. Instead, the results have to be derived via indirect means. At the heart of the proof of the convergence is a novel dimensionality argument.<sup>5</sup>

There is a great number of work on  $n$ -player non-cooperative bargaining games. Herrero (1985) and Sutton (1986) demonstrate the problems in extending the number of players beyond two. The convergence to the Nash bargaining solution typically requires that the bargaining game has a unique subgame perfect equilibrium. Various examples of such games are available in Chae and Yang (1988, 1994), Huang (2004), Krishna and Serrano (1996), and Suh and Wen (2006).

## 2 The set up

There is a set  $\{1, \dots, n\}$  of players and a nonempty, compact, convex and strictly comprehensive utility possibility set  $U \subset \mathbb{R}_+^n$ .<sup>6,7</sup> The vector of utilities is denoted by  $u = (u_1, \dots, u_n)$ , or  $u = (u_i, u_{-i})$ . The (weak and strong) *Pareto frontier* of  $U$  is then defined by  $P := \{u \in U : v \not\geq u, \text{ for all } v \in U\}$ .

Delay is costly: The present value of player  $i$ 's next period utility  $u_i$  is  $\delta^\Delta u_i$ , where  $0 < \delta < 1$  is the common discount factor, and  $\Delta > 0$  is the length between two stages.

We study a class of unanimity bargaining games, defined as follows.

- At stage  $t = 0, 1, \dots$  player  $i = (t + 1) \bmod n$  makes an offer  $v \in U$  and the players  $j \neq i$  accept or reject the offer in the ascending order of their index.
- If all  $j \neq i$  accept, then  $v$  is implemented. Otherwise the game moves to stage  $t + 1$ .

We concentrate on the *stationary* (subgame perfect Nash) equilibria of the game, where:

1. Each  $i$ 's proposal is dependent only on the continuation game.
2. Each  $i$ 's acceptance decision in period  $t$  depends only on the offer on the table and the continuation game.

---

<sup>5</sup>Kultti and Vartiainen (2007) develop a similar argument in a different context.

<sup>6</sup>Vector notation:  $x \geq y$  if  $x_i \geq y_i$  for all  $i$ ,  $x \geq y$  iff  $x \geq y$  and not  $x_i = y_i$  for all  $i$ , and  $x > y$  iff  $x_i > y_i$  for all  $i$ .

<sup>7</sup> $X \subset \mathbb{R}_+^k$  is comprehensive if  $x \in X$  and  $x \geq y \geq 0$  imply  $y \in X$ . It is strictly comprehensive if  $x \not\geq y$  for all  $x \in X - \{y\}$  implies  $x \not\geq y$  for all  $x \in X - \{y\}$ .

Define

$$\rho(i, j) = \begin{cases} i - j, & \text{if } i \geq j \\ i - j + n, & \text{if } i < j. \end{cases}$$

Given the stationarity assumption, player  $j$  always offers  $u^j$  when it is his turn to make an offer. Since player  $j$ 's offer  $v = (v_1, \dots, v_n)$  is accepted if

$$v_i \geq \delta^{\Delta\rho(i,j)} u_i^i, \text{ for all } i \neq j,$$

the equilibrium offer  $u^j$  satisfies,

$$u_i^j = \delta^{\Delta\rho(i,j)} u_i^i, \text{ for all } i, j. \quad (1)$$

Thus an equilibrium is characterized by a profile  $(u^1, \dots, u^n) \in U^n$ , where each  $u^i = (u_1^i, \dots, u_n^i) \in U$  specifies all players' payoffs when it is  $i$ 's turn to make an offer. We conclude that  $(u^1, \dots, u^n)$  is an equilibrium profile if and only if it meets (1) for all  $i, j \in \{1, \dots, n\}$ .

**Theorem 1** *A stationary equilibrium exists.*

**Proof.** Assume  $\Delta = 1$ . Let  $U^n$  be the  $n$ -copy of the utility set  $U$ . Denote a typical element of  $U^n$  by  $(u^1, \dots, u^n)$ . For any  $i, j$ , define function  $g_j^i : U \rightarrow \mathbb{R}_+$  such that

$$g_j^i(u^j) := \delta^{\rho(j,i)} \max\{u_j : (u_j, (u_k^j)_{k \neq j}) \in U\}. \quad (2)$$

By the compactness of  $U$   $g_j^i$  exists, and by the convexity of  $U$ ,  $g_j^i$  is continuous. Let  $g^i(\cdot) := (g_j^i(\cdot))_{j=1}^n : U^n \rightarrow \mathbb{R}_+^n$ . Define function  $\xi^i : U^n \rightarrow \mathbb{R}_+$  such that

$$\xi^i(u^1, \dots, u^n) := \max\{x \in \mathbb{R} : x g^i(u^1, \dots, u^n) \in U\},$$

for all  $(u^1, \dots, u^n) \in U^n$ .

By the compactness of  $U^n$ ,  $\xi^i$  is well defined, and by the convexity of  $U^n$ ,  $\xi^i$  is continuous. Construct a function  $h^i : U^n \rightarrow \mathbb{R}_+^n$  such that

$$h^i(u^1, \dots, u^n) := g^i(u^1, \dots, u^n) \min\{\xi^i(u^1, \dots, u^n), 1\},$$

for all  $(u^1, \dots, u^n) \in U^n$ .

Let  $h(u^1, \dots, u^n) = (h^1(u^1, \dots, u^n), \dots, h^n(u^1, \dots, u^n))$ . Then

$$h(u^1, \dots, u^n) : U^n \rightarrow U^n.$$

By continuity of  $(g^1, \dots, g^n)$  and  $(\xi^1, \dots, \xi^n)$   $h$  is a continuous function. By Brouwer's Theorem, there is a  $(v^1, \dots, v^n)$  in  $U^n$  such that

$$h(v^1, \dots, v^n) = (v^1, \dots, v^n). \quad (3)$$

If also

$$g(v^1, \dots, v^n) \in U^n, \quad (4)$$

then  $g(v^1, \dots, v^n) = (v^1, \dots, v^n)$ , i.e., by (2),

$$v_i^i = g_i^i(v^i) = \max\{u_i : (u_i, (v_k^i)_{k \neq i}) \in U\}, \text{ for all } i, \quad (5)$$

$$v_j^i = g_j^i(v^i) = \delta^{\rho(j,i)} \max\{u_j : (u_j, (v_k^j)_{k \neq j}) \in U\}, \text{ for all } i \neq j, \quad (6)$$

or, by plugging (5) into (6),

$$v_j^i = \delta^{\rho(j,i)} v_j^j, \text{ for all } i, j.$$

Thus (4) is sufficient for  $(v^1, \dots, v^n)$  to satisfy (1).

Suppose (4) does not hold. Then there is  $i$  such that

$$\xi^i(v^1, \dots, v^n) < 1. \quad (7)$$

By (3),

$$v^i = g^i(v^1, \dots, v^n) \xi^i(v^1, \dots, v^n).$$

Hence

$$v_j^i < g_j^i(v^1, \dots, v^n), \text{ for all } j. \quad (8)$$

By construction,

$$v^i = g^i(v^1, \dots, v^n) \xi^i(v^1, \dots, v^n) \in P.$$

Thus, by (8),

$$(g_i^i(v^i), v_{-i}^i) \notin U. \quad (9)$$

But, by (2),  $g_i^i(v^i) = \max\{u_i : (u_i, v_{-i}^i) \in U\}$ , i.e.  $(g_i^i(v^i), v_{-i}^i)$  is the element in  $U$  that maximizes  $i$ 's payoff given that the other players get at least  $v_{-i}^i$ , a contradiction to (9). ■

### 3 Relationship with the Nash solution

Denote the *Nash solution* by

$$u^N := \arg \max_{u \in U} \prod_{i=1}^n u_i. \quad (10)$$

Also denote by

$$H(u) := \left\{ (v_1, \dots, v_n) \in \mathbb{R}^n : \prod_{i=1}^n v_i = \prod_{i=1}^n u_i \right\},$$

the hyperbola that contains  $u$ . Note that, by construction,  $u^N$  is the unique point in which  $U$  is supported by a hyperbola, this time by  $H(u^N)$ .



It is not difficult to see that equilibrium offers  $u^1(\Delta), \dots, u^n(\Delta)$  under  $\Delta > 0$  lie in the same hyperbola: for any  $j$ ,

$$\prod_{i=1}^n u_i^j(\Delta) = \prod_{i=1}^n \delta^{\Delta \rho(j,i)} u_i^i(\Delta) = \delta^{\Delta n(n-1)/2} \prod_{i=1}^n u_i^i(\Delta). \quad (11)$$

The last expression is independent of the proposer index  $j$ .

**Theorem 2** *Let  $P$  be differentiable. Then all stationary equilibrium outcomes converge to  $u^N$  as  $\Delta$  tends to 0.*

The proof can be summarized as follows. Consider the three player case. Think of the surface  $P$  of  $U$  as a chart of 1-dimensional curves, each reflecting an intersection of  $P$  and a hyperbola. Identify the equilibrium offers  $u^1(\Delta), u^2(\Delta), u^3(\Delta)$  under  $\Delta$ . As  $\Delta$  becomes small, the maximum distance between vectors  $u^1(\Delta), u^2(\Delta), u^3(\Delta)$  becomes small. Since, by (11), they all lie in the same hyperbola, they must either converge to the Nash solution, or, in the limit, the vectors are contained by a 1-dimensional subspace (see Fig. 1). We show that  $u^1(\Delta), u^2(\Delta), u^3(\Delta)$  are always linearly independent, and hence cannot be embedded into a 1-dimensional subspace. Thus  $u^1(\Delta), u^2(\Delta), u^3(\Delta)$  cannot converge anywhere but to the Nash solution.

[FIGURE 1 AROUND HERE]

**Proof of Theorem 2.** Denote by  $u^j(\Delta) = (u_i^j(\Delta))_{i=1}^n$  player  $j$ 's equilibrium offer when the period length is  $\Delta > 0$ . Note that, since  $U$  is bounded, the difference  $(\delta^{-\Delta \rho(i,j)} - 1)u_i^j(\Delta)$  tends to 0 as  $\Delta$  becomes negligible for all  $i, j$ . Hence so does the difference  $u_i^i(\Delta) - u_i^j(\Delta)$ . This implies that  $u^j(\Delta)$  and  $u^i(\Delta)$ , for all  $i, j$ , approach one another as  $\Delta$  tends to 0.

Let  $\{\Delta\}$  be a subsequence under which  $u^1(\Delta), \dots, u^n(\Delta)$  converge to  $u^*$ . Since  $U$  is bounded, it suffices for us to show that  $u^* = u^N$ . All the limits below are taken with respect to the sequence  $\{\Delta\}$ .

It will be easier to work with logarithmized objects since logarithmization transforms hyperbolas into hyperplanes without affecting the nature of

local properties of the objects.<sup>8</sup> For expositional reasons, denote the logarithmized variables by

$$\begin{aligned}\bar{u} &= \ln u = (\ln u_1, \dots, \ln u_n), \text{ for any } u \in U, \\ \bar{S} &= \{\bar{u} : u \in S\}, \text{ for any } S \subseteq \mathbb{R}_{++}^n, \\ \bar{\delta} &= \ln \delta.\end{aligned}$$

Thus, by (1),

$$\bar{u}_i^j(\Delta) = \Delta \bar{\delta} \rho(i, j) + \bar{u}_i^i(\Delta), \text{ for all } i, j. \quad (12)$$

Logarithmization preserves the convexity of  $U$  and the differentiability of  $P$ . Thus  $\bar{P}$  is an  $n - 1$ -manifold supporting, at each of its point, a unique  $n - 1$  dimensional hyperplane. In particular, it supports a unique hyperplane at  $\bar{u}^*$ .

Player  $j$ 's equilibrium offer under  $\Delta > 0$  is  $\bar{u}^j(\Delta) = (\bar{u}_i^j(\Delta))_{i=1}^n$ .<sup>9</sup> Then, by (12),

$$\begin{aligned}\ln \prod_{i=1}^n u_i^j(\Delta) &= \sum_{i=1}^n \bar{u}_i^j(\Delta) \\ &= \Delta \bar{\delta} \sum_{i=1}^n \rho(i, j) + \sum_{i=1}^n \bar{u}_i^i(\Delta) \\ &= \frac{n(n-1)\Delta \bar{\delta}}{2} + \sum_{i=1}^n \bar{u}_i^i(\Delta).\end{aligned} \quad (13)$$

Construct the  $n - 1$  -dimensional hyperplane  $L^\Delta$  such that,

$$L^\Delta = \left\{ x \in \mathbb{R}^n : \sum_{i=1}^n x_i = \frac{n(n-1)\Delta \bar{\delta}}{2} + \sum_{i=1}^n \bar{u}_i^i(\Delta) \right\}.$$

Since (13) is independent of the index  $j$ , it follows that

$$\bar{u}^1(\Delta), \dots, \bar{u}^n(\Delta) \in L^\Delta. \quad (14)$$

Identify

$$C = \left\{ \left( \begin{array}{c} 0 \\ 1 \\ 2 \\ \vdots \\ n-1 \end{array} \right), \left( \begin{array}{c} n-1 \\ 0 \\ 1 \\ \vdots \\ n-2 \end{array} \right), \dots, \left( \begin{array}{c} 1 \\ 2 \\ \vdots \\ n-1 \\ 0 \end{array} \right) \right\},$$

<sup>8</sup>Since it is easier to verify whether an object can be embedded into a hyperplane than into a hyperbola.

<sup>9</sup>To be precise, logarithmic scales require that all equilibrium offers are bounded away from zero for all  $i$  and all  $\Delta > 0$ . This fact is easy to verify from the equilibrium conditions.

a collection of  $n - 1$  linearly independent vectors. By (12), the collection of equilibrium offers can now be written compactly

$$\{\bar{u}^1(\Delta), \dots, \bar{u}^n(\Delta)\} = (\bar{u}_i^i(\Delta))_{i=1}^n + \Delta\bar{\delta}C. \quad (15)$$

Thus, by (14),

$$(\bar{u}_i^i(\Delta))_{i=1}^n + \Delta\bar{\delta}C \subset L^\Delta, \quad \text{for all } \Delta.$$

By making an affine transformation of both sides,<sup>10</sup>

$$C \subset \frac{1}{\Delta\bar{\delta}}L^\Delta - \frac{1}{\Delta\bar{\delta}}(\bar{u}_i^i(\Delta))_{i=1}^n, \quad \text{for all } \Delta. \quad (16)$$

Let  $L(\bar{u})$  be the hyperplane that supports  $\bar{U}$  at  $\bar{u}$ .<sup>11</sup> Since  $P$  is compact and differentiable, there is a function  $\sigma > 0$  such that  $\sigma(\varepsilon) \rightarrow 0$  as  $\varepsilon \rightarrow 0$ , and such that (see Fig. 2)

$$\sigma(\|\bar{u} - \bar{v}\|) \geq \frac{\min\{\|\bar{v} - \bar{v}'\| : \bar{v}' \in L(\bar{u})\}}{\|\bar{u} - \bar{v}\|}, \quad \text{for all } \bar{u}, \bar{v} \in \bar{P}. \quad (17)$$

For a formal proof of this, see Proposition 3 in the appendix.

[FIGURE 2 AROUND HERE]

By (17),

$$\sigma(\|\bar{u}^i(\Delta) - \bar{u}^1(\Delta)\|) \geq \frac{\min\{\|\bar{u}^i(\Delta) - \bar{v}\| : \bar{v} \in L(\bar{u}^1(\Delta))\}}{\|\bar{u}^i(\Delta) - \bar{u}^1(\Delta)\|}. \quad (18)$$

Denote by

$$\sigma^\Delta = \max_{i=1, \dots, n} \sigma(\|\bar{u}^i(\Delta) - \bar{u}^1(\Delta)\|).$$

Since  $\|\bar{u}^i(\Delta) - \bar{u}^1(\Delta)\| \rightarrow 0$  for all  $i$  it follows that  $\sigma^\Delta \rightarrow 0$  as  $\Delta \rightarrow 0$ .

Let  $\|X\|$  be the sup-norm of the set  $X$ , i.e.  $\|X\| = \sup_{x, y \in X} \|x - y\|$ . By (15),

$$\|\bar{u}^i(\Delta) - \bar{u}^1(\Delta)\| \leq \Delta\bar{\delta}\|C\|, \quad \text{for all } i. \quad (19)$$

<sup>10</sup>By an affine transformation of a set  $X \subset \mathbb{R}^m$  we mean a map  $X \mapsto aX + b$ , for  $a \in \mathbb{R}_{++}$  and  $b \in \mathbb{R}^m$ , where  $aX + b = \{ax + b : x \in X\}$ .

<sup>11</sup>That is,  $L(\bar{u}) = \{x \in \mathbb{R}^n : px = k\}$ , where  $p \in \mathbb{R}^n$  and  $k \in \mathbb{R}$  such that  $p\bar{v} \geq k$ , for all  $\bar{v} \in \bar{U}$ , and  $p\bar{u} = k$ .

By (18) and (19),

$$\min\{\|\bar{u}^i(\Delta) - \bar{v}\| : \bar{v} \in L(\bar{u}^1(\Delta))\} \leq \sigma^\Delta \Delta \bar{\delta} \|C\|, \quad \text{for all } i. \quad (20)$$

From (20) we have that

$$(\bar{u}_i^i(\Delta))_{i=1}^n + \Delta \bar{\delta} C \subset \{\bar{u} : \|\bar{u} - \bar{v}\| < \sigma^\Delta \Delta \bar{\delta} \|C\| \text{ and } \bar{v} \in L(\bar{u}^1(\Delta))\}.$$

Or, by making an affine transformation of both sides,

$$\begin{aligned} C &\subset \frac{1}{\bar{\delta}\Delta} [\{\bar{u} : \|\bar{u} - \bar{v}\| < \sigma^\Delta \Delta \bar{\delta} \|C\| \text{ and } \bar{v} \in L(\bar{u}^1(\Delta))\} - (\bar{u}_i^i(\Delta))_{i=1}^n] \\ &= \left\{ \bar{u} : \|\bar{u} - \bar{v}\| < \sigma^\Delta \|C\| \text{ and } \bar{v} \in \frac{1}{\bar{\delta}\Delta} L(\bar{u}^1(\Delta)) \right\} - \frac{1}{\bar{\delta}\Delta} (\bar{u}_i^i(\Delta))_{i=1}^n. \end{aligned} \quad (21)$$

That is,  $C$  is contained in the  $\sigma^\Delta \|C\|$ -neighborhood of an  $n-1$ -dimensional hyperplane  $(\bar{\delta}\Delta)^{-1} L(\bar{u}^1(\Delta)) - (\bar{\delta}\Delta)^{-1} (\bar{u}_i^i(\Delta))_{i=1}^n$ .

Define

$$L^* = \left\{ x \in \mathbb{R}^n : \sum_{i=1}^n x_i = \sum_{i=1}^n \bar{u}_i^* \right\}.$$

By construction,  $\bar{u}^* \in L^*$ . Suppose that  $L(\bar{u}^*) \neq L^*$ , i.e. the hyperplane that supports  $\bar{U}$  at  $\bar{u}^*$  does not coincide with the hyperplane  $L^*$ . Since both  $L(\bar{u}^*)$  and  $L^*$  are  $n-1$  dimensional hyperplanes, the hyperplane  $L(\bar{u}^*) \cap L^*$  is only  $n-2$  dimensional. Since  $C$  contains  $n-1$  independent vectors, and  $\sigma^\Delta \rightarrow 0$ , there is  $\Delta' > 0$  such that for all  $\Delta < \Delta'$ ,

$$C \not\subset \left\{ \bar{u} : \|\bar{u} - \bar{v}\| < \sigma^\Delta \|C\| \text{ and } \bar{v} \in \frac{1}{\bar{\delta}\Delta} L(\bar{u}^*) \right\} \cap \frac{1}{\bar{\delta}\Delta} L^* - \frac{1}{\bar{\delta}\Delta} \bar{u}^*.$$

Since  $\bar{P}$  is differentiable and  $\bar{u}^1(\Delta) \rightarrow \bar{u}^*$ , it follows that  $L^\Delta \rightarrow L^*$  and  $L(\bar{u}^1(\Delta)) \rightarrow L(\bar{u}^*)$ .<sup>12</sup> Thus there is  $\Delta'' < \Delta'$  such that for all  $\Delta < \Delta''$ ,

$$C \not\subset \left\{ \bar{u} : \|\bar{u} - \bar{v}\| < \sigma^\Delta \|C\| \text{ and } \bar{v} \in \frac{1}{\bar{\delta}\Delta} L(\bar{u}^1(\Delta)) \right\} \cap \frac{1}{\bar{\delta}\Delta} L^\Delta - \frac{1}{\bar{\delta}\Delta} (\bar{u}_i^i(\Delta))_{i=1}^n,$$

which contradicts (16) and (21). Thus  $L(\bar{u}^*) = L^*$ .

Since  $L(\bar{u}^*) = L^*$ ,  $L^*$  supports  $\bar{U}$  at  $\bar{u}^*$ . Since, by definition,  $L^*$  is the log-transformed hyperbola  $H(u^*)$ , and  $\bar{U}$  is the log-transformed utility set  $U$ , this must mean that the  $U$  is supported by the hyperbola  $H(u^*)$  at  $u^*$ . Thus  $u^* = u^N$ . ■

<sup>12</sup>Where the limits are defined with respect to the normals of the hyperplanes.

## 4 Necessity of the Differentiability of $P$

The result of the previous section, which is the main finding of the paper, is based on the local properties of the Pareto frontier, i.e. it shows that the Nash program works in any environment where the Pareto frontier is locally isomorphic to a hyperplane. However, when this is not the case, the result need not hold. Indeed, to this end we argue that the convergence result is sensitive to the differentiability assumption. We give an example of a scenario where stationary equilibria do not converge to the Nash bargaining solution.

Let

$$U = \{u \in \mathbb{R}_+^3 : u_1 + u_2 \leq 1, u_3 \leq 1\}.$$

Here 1 and 2 bargain over a linear cake and 3 is a "dummy" player, without strategic significance.

Equilibrium condition (1) implies that the stationary equilibrium offers  $u^1, u^2$ , and  $u^3$  satisfy

$$\begin{aligned} \delta^{2\Delta} u_1^1 &= \delta^\Delta u_1^3 = u_1^2, \\ \delta^{2\Delta} u_2^2 &= \delta^\Delta u_2^1 = u_2^3, \\ \delta^{2\Delta} u_3^3 &= \delta^\Delta u_3^2 = u_3^1. \end{aligned} \tag{22}$$

In equilibrium, players do not waste their own consumption possibilities when making offers. This means

$$\begin{aligned} u_1^1 &= 1 - u_2^1, \\ u_2^2 &= 1 - u_1^2, \\ u_3^3 &= 1. \end{aligned} \tag{23}$$

Combining (23) with (22) gives the equilibrium offers for players 1, 2 and 3,

$$\begin{aligned} u^1 &= \left( \frac{1 - \delta^\Delta}{1 - \delta^{3\Delta}}, \frac{\delta^\Delta - \delta^{3\Delta}}{1 - \delta^{3\Delta}}, \delta^{2\Delta} \right), \\ u^2 &= \left( \frac{\delta^{2\Delta} - \delta^{3\Delta}}{1 - \delta^{3\Delta}}, \frac{1 - \delta^{2\Delta}}{1 - \delta^{3\Delta}}, \delta^\Delta \right), \\ u^3 &= \left( \frac{\delta^\Delta - \delta^{2\Delta}}{1 - \delta^{3\Delta}}, \frac{\delta^{2\Delta} - \delta^{4\Delta}}{1 - \delta^{3\Delta}}, 1 \right). \end{aligned}$$

By taking the limit  $\Delta \rightarrow 0$ , we obtain the common convergence point of  $u^1, u^2$ , and  $u^3$

$$u^* = \left( \frac{1}{3}, \frac{2}{3}, 1 \right).$$

However, by symmetry, the Nash solution of the problem is

$$u^N = \left( \frac{1}{2}, \frac{1}{2}, 1 \right).$$

Thus the convergence point  $u^*$  of the stationary equilibrium does not coincide with the Nash solution  $u^N$  (see Fig. 3). Hence the differentiability of  $P$  is crucial for the convergence result (the seminal idea is by Lensberg and Thomson, 1988).

[FIGURE 3 AROUND HERE]

However, our equilibrium is not the only stationary equilibrium that  $U$  entertains. This is due to the fact that  $U$  is comprehensive but not strictly comprehensive, as demanded by our characterization (1). However, using the above reasoning, it is clear that  $U$  can be approximated by a strictly comprehensive problem whose unique stationary equilibrium is close to the equilibrium we characterize.<sup>13</sup> Thus the no-convergence result does hold also in the class of convex, compact, and strictly comprehensive problems.

**Discussion** To understand why the convergence to the Nash solution does not always hold, note that in the above example, there is no tradeoff between player 3 and the other players. Players 1 and 2 effectively bargain against one another, and the only effect of the existence of 3 is to delay 1's offer once 2 has rejected his offer. The role of differentiability is to guarantee that the Pareto surface is locally fully competitive; a small chunk of payoff of one player can be distributed to others in any proportion. This means that the details of the players' intra game relations, e.g. dummyness as in the above example, do not affect bargaining when the discount factor becomes small. Hence, in the limit, each player regards the other players symmetrically. Because of the convexity of the problem, this property is met locally only in the Nash solution. This is what drives the convergence result.

In the two player context, however, there is no problem in treating all opponents symmetrically as each player only faces one opponent. This guarantees convergence even when the Pareto surface is not differentiable. Technically, this is mirrored by the fact that the intersection of the Pareto surface and a hyperbola is zero dimensional, and hence cannot be connected. Because of this, the shrinking set of players equilibrium offers cannot converge

---

<sup>13</sup>Think of the problem  $U^\varepsilon = U(1 - \varepsilon) + V\varepsilon$ , where  $V = \{u \in \mathbb{R}_+^3 : 2u_1 + 2u_2 + u_3 \leq 3\}$ , and  $\varepsilon < 0$ . Now  $U^\varepsilon$  is strictly comprehensive and for small  $\varepsilon$ , the unique stationary equilibrium converges to a point close to  $u^*$ .

anywhere but to the point of intersection of the highest hyperbola and the Pareto surface.

To interpret this observation in terms of the so called "Nash program", which aims at reconciling the strategic approach and the axiomatic one, note that a non-differentiable utility space can always be approximated by differentiable ones. Since the Nash solution is continuous with respect to small changes in the Pareto surface, it must be the case that the noncooperative equilibrium correspondence is discontinuous with respect to the underlying payoff parameters. Thus the "problem" seems to be with the noncooperative approach; (the limit) outcome of noncooperative bargaining can be sensitive to the fine details of the underlying physical model.

On the other hand, the non-convergence result may be interpreted as a manifestation of the fact that the Nash IIA axiom does not adequately take into account categorial changes in players' relationships, e.g. that one player becomes completely independent in terms of payoffs. Indeed, replacing IIA with the multilateral stability condition of Lensberg and Thomson (1988) (see also Thomson and Lensberg, 1990, ch 8), whose target is precisely that, is known to characterize the Nash solution only in the domain of smooth problems.

## References

- [1] BINMORE K., A. RUBINSTEIN AND A. WOLINSKY 1986, The Nash bargaining solution in economic modelling, *Rand Journal of Economics* 17, 176-188.
- [2] BINMORE K. 1985, Bargaining and coalitions, in Roth A. (ed.): *Game Theoretic Models of Bargaining*. New York: Cambridge University Press.
- [3] CHAE S. AND J.-A. YANG 1988, The unique perfect equilibrium of an N-person bargaining game, *Economics Letters* 28, 221-223.
- [4] CHAE S. AND J.-A. YANG 1994, An N-person pure bargaining game, *Journal of Economic Theory* 62, 86-102.
- [5] CHATTERJEE, K. AND H. SABOURIAN 2000, Multiperson bargaining and strategic complexity, *Econometrica* 68, 1491-1509
- [6] HERRERO, M. 1985, Strategic theory of market institutions, unpublished Ph.D dissertation, LSE.
- [7] HUANG C.-Y. 2002, Multilateral bargaining: conditional and unconditional offers, *Economic Theory* 20, 401-412

- [8] KRISHNA V. AND R. SERRANO 1996, Multilateral bargaining, *Review of Economic Studies* 63, 61-80.
- [9] KULTTI, K. AND H. VARTAINEN 2005, Von Neumann-Morgenstern stable sets, discounting, and Nash bargaining, *Journal of Economic Theory*, *forthcoming*.
- [10] LENSBERG, T. AND W. THOMSON 1988, Characterizing the Nash solution without Parato-optimality, *Social Choice and Welfare* 5, 547-59.
- [11] NASH J. 1950, The bargaining problem, *Econometrica* 18, 155-162.
- [12] RUBINSTEIN A. 1982, Perfect equilibrium in a bargaining model, *Econometrica* 50, 97-109.
- [13] SUH S.-C. AND Q. WEN 2006, Multi-agent bilateral bargaining and the Nash bargaining solution, *Journal of Mathematical Economics* 42, 61-73.
- [14] SUTTON J. 1986, Non-cooperative bargaining theory: an introduction, *Review of Economic Studies* 53,709-724.
- [15] THOMSON, W. AND T. LENSBERG 1989, Axiomatic theory of bargaining with variable number of agents, Cambridge UK: Cambridge University Press.
- [16] VON NEUMANN J. AND O. MORGENSTERN 1944, *Theory of Games and Economic Behavior*. New York: John Wiley and Sons.



## A Appendix

In the following proposition, all variables are defined as in Theorem 2.

**Proposition 3** *There is a function  $\sigma > 0$  such that  $\sigma(\varepsilon) \rightarrow 0$  as  $\varepsilon \rightarrow 0$  and such that*

$$\sigma(\|\bar{u} - \bar{v}\|) \geq \frac{\min\{\|\bar{v} - \bar{v}'\| : \bar{v}' \in L(\bar{u})\}}{\|\bar{u} - \bar{v}\|}, \text{ for all } \bar{u}, \bar{v} \in \bar{P}.$$

**Proof.** Since  $\bar{P}$  is differentiable, it supports at every  $\bar{u} \in \bar{P}$  an  $n$ -dimensional ball with radius  $r(\bar{u}) > 0$ . Since  $\bar{P}$  is differentiable, and a ball is locally isomorphic to a hyperplane, it suffices for us to show that  $r$  is uniformly bounded below by some  $\mu > 0$  on  $\bar{P}$ .

If  $r$  is not uniformly bounded by any  $\mu > 0$  on  $\bar{P}$ , then there is a sequence  $\{\bar{u}^k\}$  of elements in  $\bar{P}$  such that  $r(\bar{u}^k)$  converges to 0. Since  $\bar{P}$  is differentiable at  $\bar{u}$  if and only if  $P$  is differentiable at  $u$ , it follows that there is a corresponding sequence  $\{u^k\}$  of elements in  $P$  such that  $r(u^k)$ , the strictly positive radius of a ball supporting  $P$  at  $u^k$ , converges to 0. But this cannot hold since  $P$  is compact and differentiable. ■

Figures




Figure 1




Figure 2




Figure 3

**Aboa Centre for Economics (ACE)** was founded in 1998 by the departments of economics at the Turku School of Economics, Åbo Akademi University and University of Turku. The aim of the Centre is to coordinate research and education related to economics in the three universities.

Contact information: Aboa Centre for Economics, Turku School of Economics, Rehtorinpellonkatu 3, 20500 Turku, Finland.

**Aboa Centre for Economics (ACE)** on Turun kolmen yliopiston vuonna 1998 perustama yhteistyöelin. Sen osapuolet ovat Turun kauppakorkeakoulun kansantaloustieteen oppiaine, Åbo Akademin national-ekonomi-oppiaine ja Turun yliopiston taloustieteen laitos. ACEn toiminta-ajatuksena on koordinoida kansantaloustieteen tutkimusta ja opetusta Turun kolmessa yliopistossa.

Yhteystiedot: Aboa Centre for Economics, Kansantaloustiede, Turun kauppakorkeakoulu, 20500 Turku.

[www.tse.fi/ace](http://www.tse.fi/ace)

ISSN 1796-3133